

CP-1228

PATENT APPLICATION  
IN THE  
UNITED STATES PATENT AND TRADEMARK OFFICE

TITLE: PRIMARY INTERMEDIATES FOR  
OXIDATIVE COLORATION OF HAIR

Inventors: Mu-III Lim  
Yuh-Guo Pan

10052741.011802

## PRIMARY INTERMEDIATES FOR OXIDATIVE COLORATION OF HAIR

### Field of the Invention

5

This invention relates to new 2-substituted 4-amino-phenol derivative compounds and compositions containing these compounds, as primary intermediates for oxidative coloring of hair fibers.

10

### Background to the Invention

15

Coloration of hair is a procedure practiced from antiquity employing a variety of means. In modern times, the method most extensively to color hair is an oxidative dyeing process utilizing one or more oxidative hair coloring agents in combination with one or more oxidizing agents.

20

Most commonly a peroxy oxidizing agent is used in combination with one or more oxidative hair coloring agents, generally small molecules capable of diffusing into hair and comprising one or more primary intermediates and one or more couplers. In this procedure, a peroxide material, such as hydrogen peroxide, is employed to activate the small molecules of primary intermediates so that they react with couplers to form larger sized compounds in the hair shaft to color the hair in a variety of shades and colors.

25

A wide variety of primary intermediates and couplers have been employed in such oxidative hair coloring systems and compositions. Among the primary intermediates employed there may be mentioned p-phenylenediamine, p-toluenediamine, p-aminophenol, 4-amino-3-methylphenol, N,N-bis(2-hydroxyethyl)-p-phenylene diamine, 1-(2-hydroxyethyl)-4,5-diaminopyrazole and as couplers there may be mentioned

30

resorcinol, 2-methylresorcinol, 3-aminophenol, 2,4-diaminophenoxyethanol, and 5-amino-2-methylphenol.

There are numerous additional requirements for oxidation dye compounds that are used to dye human hair besides the color or the desired intensity. Thus, the dye compounds must be unobjectionable in regard to toxicological and dermatological properties and must provide the desired hair color with a good light fastness, fastness to a permanent wave treatment, acid fastness and fastness to rubbing. The color of the hair dyed with the dye compounds in each case must be stable for at least 4 to 6 weeks to light, rubbing and chemical agents. Furthermore, an additional requirement is the production of a broad palette of different color shades using different developer and coupler substances. Many of the desired shades have been produced with dyes based on p-aminophenol. However, as indicated in U. S. Patent No. 4,997,451, the use of p-aminophenol is being questioned for possible toxicological reasons. The proposed replacements for p-aminophenol have not proved entirely satisfactory. There is therefore a need for new primary intermediate compounds to meet one or more of the desired properties but not possessing the possible toxicological drawbacks possessed by p-aminophenol.

### Summary of the Invention


It is therefore an object of this invention to provide new primary intermediate compounds useful in place of p-aminophenol to provide a wide range of different color shades with various combinations of primary intermediates and couplers, but which avoids the drawbacks of p-aminophenol.

It has been discovered that new certain 2-substituted 4-amino-phenol derivative compounds are suitable primary intermediates for hair coloring compositions and systems for providing good oxidative coloration of

hair and for providing acceptable light fastness, good selectivity, fastness to shampooing, fastness to perspiration and permanent wave treatment, and suitable for providing a wide variety of different color shades with various primary intermediate and coupler compounds.

5

The invention provides new compounds of Formula (1):




(1)

wherein  $R_1$  and  $R_2$  are each individually selected from hydrogen atoms,  $C_1$  to  $C_5$  alkyl,  $C_1$  to  $C_5$  mono or dihydroxyalkyl, phenyl or benzyl optionally substituted with a hydroxyl, amino or  $C_1$  to  $C_3$  alkoxy group, or  $R_1$  and  $R_2$  together with the nitrogen atom to which they are attached form a  $C_3$  to  $C_6$ , preferably  $C_4$  to  $C_6$ , saturated or unsaturated ring optionally containing in the ring one or more additional hetero atoms selected from O, S and N atoms.

These novel primary intermediates are used to provide coloration to hair in which there is good dye uptake by the hair and provides shades or colors which are stable over a relatively long period of time. The novel primary intermediates provide for dyeing of hair to impart color or shades that possess good wash fastness and do not undergo significant changes on exposure to light or shampooing.

## Detailed Description of the Invention

Preferred 4-amino-phenol derivative compounds of this invention are those of formula (1)




5 wherein  $R_1$  and  $R_2$  are each individually selected from a hydrogen atom, a  $C_1$  to  $C_3$  alkyl group, preferably methyl groups; phenyl or benzyl optionally substituted with an alkoxy group, preferably a methoxy group, or  $R_1$  and  $R_2$  together with the nitrogen atom to which they are bound form a piperazine, piperidine, imidazole, or morpholine ring.

Especially preferred compounds of formula (1) of this invention are the following compounds:

- 4-amino-2-(1-phenylamino-ethyl)-phenol;
- 15 4-amino-2-(1-piperidin-1-yl-ethyl)-phenol;
- 4-amino-2-[1-(pyridin-3-yl-amino)-ethyl]-phenol;
- 4-amino-2-(1-benzylamino-ethyl)phenol;
- 4-amino-2-[1-(4-methoxy)-phenylamino-ethyl]-phenol;
- 4-amino-2-(1-amino-ethyl)-phenol;
- 20 4-amino-2-(1-dimethylamino-ethyl)-phenol;
- 4-amino-2-(1-morpholin-4-yl-ethyl)-phenol;
- 4-amino-2-[1-(4-methoxy)-benzylamino-ethyl]-phenol;
- 4-amino-2-(1-methylamino-ethyl)-phenol; and
- 4-amino-2-[1-(2-hydroxyethylamino)-ethyl]-phenol.

The new compounds of formula (1) of this invention can be prepared according to the following reaction sequence wherein  $R_1$  and  $R_2$  are as defined hereinbefore:



In this synthesis procedure 5-butylamido-2-hydroxyacetophenone of formula (2) is subjected to reductive amination with the reagent  $R_1R_2NH$  using reducing agents such as sodium triacetoxyborohydride, sodium cyanoborohydride or polymethylhydrosiloxane (PMHS) to produce a compound of formula (3). Hydrolysis of the compound of formula (3) with methanolic hydrochloric acid produces a compound of formula (1).

### Synthesis Examples 1 to 11

Employing the appropriate  $R_1R_2NH$  reagent, reducing agent and 5-butylamido-2-hydroxyacetophenone in the synthesis procedure described, the following compounds of formula (1) of this invention can be prepared:

- 4-amino-2-(1-phenylamino-ethyl)-phenol;
- 4-amino-2-(1-piperidin-1-yl-ethyl)-phenol;
- 4-amino-2-[1-(pyridin-3-yl-amino)-ethyl]-phenol;
- 4-amino-2-(1-benzylamino-ethyl)-phenol;
- 4-amino-2-[1-(4-methoxy)-phenylamino-ethyl]-phenol;
- 4-amino-2-(1-amino-ethyl)-phenol;

4-amino-2-(1-dimethylamino-ethyl)-phenol;  
 4-amino-2-(1-morpholin-4-yl-ethyl)-phenol;  
 4-amino-2-[1-(4-methoxy)-benzylamino-ethyl]-phenol;  
 4-amino-2-(1-methylamino-ethyl)-phenol; and  
 4-amino-2-[1-(2-hydroxyethylamino)-ethyl]-phenol.

As used herein, the term "hair dyeing composition" (also  
 synonymously referred to herein as the hair dye composition, the hair coloring  
 composition, or the hair dye lotion) refers to the composition containing  
 oxidation dyes, including the novel compounds described herein, prior to  
 admixture with the developer composition. The term "developer composition"  
 (also referred to as the oxidizing agent composition or the peroxide  
 composition) refers to compositions containing an oxidizing agent prior to  
 admixture with the hair dyeing composition. The term "hair dye product" or  
 "hair dye system" (also referred to as the hair dyeing system, hair dyeing  
 product, or hair coloring system) interchangeably refer to the combination of  
 the hair dyeing composition and the developer composition before admixture,  
 and may further include a conditioner product and instructions, such product  
 or system often being provided packaged as a kit. The term "hair dyeing  
 product composition" refers to the composition formed by mixing the hair  
 dyeing composition and the developer composition. "Carrier" (or vehicle or  
 base) refers to the combination of ingredients contained in a composition  
 excluding the active agents (e.g., the oxidation hair dyes of the hair dyeing  
 composition).

25

Hair coloring (i.e., hair dyeing) compositions of this invention  
 can contain, in combination with oxidation dye couplers, a novel primary  
 intermediate of this invention as the sole primary intermediate or can also  
 contain other primary intermediates. Thus, one or more suitable primary  
 intermediates may be used in combination with the novel primary  
 intermediates of this invention.

Suitable known primary intermediates include, for example,

p-phenylenediamine derivatives such as: benzene-1,4-diamine (commonly known as p-phenylenediamine), 2-methyl-benzene-1,4-diamine, 2-chloro-benzene-1,4-diamine, N-phenyl-benzene-1,4-diamine, N-(2-ethoxyethyl)benzene-1,4-diamine, 2-[(4-amino-phenyl)-(2-hydroxy-ethyl)-amino]-ethanol, (commonly known as N,N-bis(2-hydroxyethyl)-p-phenylenediamine) (2,5-diamino-phenyl)-methanol, 1-(2,5-diamino-phenyl)-ethanol, 2-(2,5-diamino-phenyl)-ethanol, N-(4-aminophenyl)benzene-1,4-diamine, 2,6-dimethyl-benzene-1,4-diamine, 2-isopropyl-benzene-1,4-diamine, 1-[(4-aminophenyl)amino]-propan-2-ol, 2-propyl-benzene-1,4-diamine, 1,3-bis[(4-aminophenyl)(2-hydroxyethyl)amino]propan-2-ol, N<sup>4</sup>,N<sup>4</sup>,2-trimethylbenzene-1,4-diamine, 2-methoxy-benzene-1,4-diamine, 1-(2,5-diaminophenyl)ethane-1,2-diol, 2,3-dimethyl-benzene-1,4-diamine, N-(4-amino-3-hydroxy-phenyl)-acetamide, 2,6-diethylbenzene-1,4-diamine, 2,5-dimethylbenzene-1,4-diamine, 2-thien-2-ylbenzene-1,4-diamine, 2-thien-3-ylbenzene-1,4-diamine, 2-pyridin-3-ylbenzene-1,4-diamine, 1,1'-biphenyl-2,5-diamine, 2-(methoxymethyl)benzene-1,4-diamine, 2-(aminomethyl)benzene-1,4-diamine, 2-(2,5-diaminophenoxy)ethanol, N-[2-(2,5-diaminophenoxy)ethyl]-acetamide, N,N-dimethylbenzene-1,4-diamine, N,N-diethylbenzene-1,4-diamine, N,N-dipropylbenzene-1,4-diamine, 2-[(4-aminophenyl)(ethyl)amino]ethanol, 2-[(4-amino-3-methyl-phenyl)-(2-hydroxy-ethyl)-amino]-ethanol, N-(2-methoxyethyl)-benzene-1,4-diamine, 3-[(4-aminophenyl)amino]propan-1-ol, 3-[(4-aminophenyl)-amino]propane-1,2-diol, N-{4-[(4-aminophenyl)amino]butyl}benzene-1,4-diamine, and 2-[2-(2-{2-[(2,5-diaminophenyl)-oxy]ethoxy}ethoxy)ethoxy]benzene-1,4-diamine;

p-aminophenol derivatives such as: 4-amino-phenol (commonly known as p-aminophenol), 4-methylamino-phenol, 4-amino-3-methyl-phenol, 4-amino-2-hydroxymethyl-phenol, 4-amino-2-methyl-phenol, 4-amino-2-[(2-hydroxy-ethylamino)-methyl]-phenol, 4-amino-2-methoxymethyl-phenol, 5-amino-2-hydroxy-benzoic acid, 1-(5-amino-2-hydroxy-phenyl)-ethane-1,2-diol,



4-amino-2-(2-hydroxy-ethyl)-phenol, 4-amino-3-(hydroxymethyl)phenol, 4-amino-3-fluoro-phenol, 4-amino-2-(aminomethyl)-phenol, and 4-amino-2-fluoro-phenol;

5 o-aminophenol derivatives such as: 2-amino-phenol (commonly known as o-aminophenol), 2,4-diaminophenol, 2-amino-5-methyl-phenol, 2-amino-6-methyl-phenol, N-(4-amino-3-hydroxy-phenyl)-acetamide, and 2-amino-4-methyl-phenol; and

10 heterocyclic derivatives such as: pyrimidine-2,4,5,6-tetramine (commonly known as 2,4,5,6-tetraaminopyridine), 1-methyl-1H-pyrazole-4,5-diamine, 2-(4,5-diamino-1H-pyrazol-1-yl)ethanol, N<sup>2</sup>,N<sup>2</sup>-dimethyl-pyridine-2,5-diamine, 2-[(3-amino-6-methoxypyridin-2-yl)amino]ethanol, 6-methoxy-N<sup>2</sup>-methyl-pyridine-2,3-diamine, 2,5,6-triaminopyrimidin-4(1H)-one, pyridine-2,5-diamine,  
15 1-isopropyl-1H-pyrazole-4,5-diamine, 1-(4-methylbenzyl)-1H-pyrazole-4,5-diamine, 1-(benzyl)-1H-pyrazole-4,5-diamine, and 1-(4-chlorobenzyl)-1H-pyrazole-4,5-diamine.

20 The novel primary intermediates of formula (1) of this invention may be used with any suitable coupler(s) in hair coloring compositions or systems of this invention.

Suitable known couplers include, for example:

25 phenols, resorcinol and naphthol derivatives such as: naphthalene-1,7-diol, benzene-1,3-diol, 4-chlorobenzene-1,3-diol, naphthalen-1-ol, 2-methyl-naphthalen-1-ol, naphthalene-1,5-diol, naphthalene-2,7-diol, benzene-1,4-diol, 2-methyl-benzene-1,3-diol, 7-amino-4-hydroxy-naphthalene-2-sulfonic acid, 2-isopropyl-5-methylphenol, 1,2,3,4-tetrahydro-naphthalene-1,5-diol, 2-chloro-benzene-1,3-diol, 4-hydroxy-naphthalene-1-sulfonic acid, benzene-1,2,3-triol, naphthalene-2,3-diol, 5-dichloro-2-

30

methylbenzene-1,3-diol, 4,6-dichlorobenzene-1,3-diol, and 2,3-dihydroxy-[1,4]naphthoquinone;

m-phenylenediamines such as: 2,4-diaminophenol, benzene-  
 5 1,3-diamine, 2-(2,4-diamino-phenoxy)-ethanol, 2-[(3-amino-phenyl)-(2-hydroxy-ethyl)-amino]-ethanol, 2-methyl-benzene-1,3-diamine, 2-[[2-(2,4-diamino-phenoxy)-ethyl]-(2-hydroxy-ethyl)-amino]-ethanol, 4-{3-[(2,4-diaminophenyl)oxy]propoxy}benzene-1,3-diamine, 2-(2,4-diamino-phenyl)-ethanol, 2-(3-amino-4-methoxy-phenylamino)-ethanol, 4-(2-amino-ethoxy)-  
 10 benzene-1,3-diamine, (2,4-diamino-phenoxy)-acetic acid, 2-[2,4-diamino-5-(2-hydroxy-ethoxy)-phenoxy]-ethanol, 4-ethoxy-6-methyl-benzene-1,3-diamine, 2-(2,4-diamino-5-methyl-phenoxy)-ethanol, 4,6-dimethoxy-benzene-1,3-diamine, 2-[3-(2-hydroxy-ethylamino)-2-methyl-phenylamino]-ethanol, 3-(2,4-diamino-phenoxy)-propan-1-ol, N-[3-(dimethylamino)phenyl]urea, 4-methoxy-  
 15 6-methylbenzene-1,3-diamine, 4-fluoro-6-methylbenzene-1,3-diamine, 2-[(3-[(2-hydroxyethyl)amino]-4,6-dimethoxyphenyl)-amino]ethanol, 3-(2,4-diaminophenoxy)-propane-1,2-diol, 2-[2-amino-4-(methylamino)-phenoxy]ethanol, 2-[(5-amino-2-ethoxy-phenyl)-(2-hydroxy-ethyl)-amino]-ethanol, 2-[(3-aminophenyl)amino]ethanol, N-(2-aminoethyl)benzene-1,3-  
 20 diamine, 4-[[2,4-diamino-phenyl)oxy]methoxy}-benzene-1,3-diamine, and 2,4-dimethoxybenzene-1,3-diamine;

m-aminophenols such as: 3-amino-phenol, 2-(3-hydroxy-4-methyl-phenylamino)-acetamide, 2-(3-hydroxy-phenylamino)-acetamide, 5-  
 25 amino-2-methyl-phenol, 5-(2-hydroxy-ethylamino)-2-methyl-phenol, 5-amino-2,4-dichloro-phenol, 3-amino-2-methyl-phenol, 3-amino-2-chloro-6-methyl-phenol, 5-amino-2-(2-hydroxy-ethoxy)-phenol, 2-chloro-5-(2,2,2-trifluoroethylamino)-phenol, 5-amino-4-chloro-2-methyl-phenol, 3-cyclopentylamino-phenol, 5-[(2-hydroxyethyl)amino]-4-methoxy-2-methylphenol, 5-amino-4-  
 30 methoxy-2-methylphenol, 3-(dimethylamino)phenol, 3-(diethylamino)phenol, 5-amino-4-fluoro-2-methylphenol, 5-amino-4-ethoxy-2-methylphenol, 3-amino-2,4-dichloro-phenol, 3-[(2-methoxyethyl)amino]phenol, 3-[(2-

hydroxyethyl)amino]phenol, 5-amino-2-ethyl-phenol, 5-amino-2-methoxyphenol, 5-[(3-hydroxypropyl)amino]-2-methylphenol, 3-[(3-hydroxy-2-methylphenyl)-amino]propane-1,2-diol, and 3-[(2-hydroxyethyl)amino]-2-methylphenol; and

5

heterocyclic derivatives such as: 3,4-dihydro-2H-1,4-benzoxazin-6-ol, 4-methyl-2-phenyl-2,4-dihydro-3H-pyrazol-3-one, 6-methoxyquinolin-8-amine, 4-methylpyridine-2,6-diol, 2,3-dihydro-1,4-benzodioxin-5-ol, 1,3-benzodioxol-5-ol, 2-(1,3-benzodioxol-5-ylamino)ethanol, 3,4-dimethylpyridine-2,6-diol, 5-chloropyridine-2,3-diol, 2,6-dimethoxypyridine-3,5-diamine, 1,3-benzodioxol-5-amine, 2-[[3,5-diamino-6-(2-hydroxy-ethoxy)-pyridin-2-yl]oxy]-ethanol, 1H-indol-4-ol, 5-amino-2,6-dimethoxypyridin-3-ol, 1H-indole-5,6-diol, 1H-indol-7-ol, 1H-indol-5-ol, 1H-indol-6-ol, 6-bromo-1,3-benzodioxol-5-ol, 2-aminopyridin-3-ol, pyridine-2,6-diamine, 3-[(3,5-diaminopyridin-2-yl)oxy]propane-1,2-diol, 5-[(3,5-diaminopyridin-2-yl)oxy]pentane-1,3-diol, 1H-indole-2,3-dione, indoline-5,6-diol, 3,5-dimethoxypyridine-2,6-diamine, 6-methoxypyridine-2,3-diamine, and 3,4-dihydro-2H-1,4-benzoxazin-6-amine.

20

Preferred primary intermediates include:

p-phenylenediamine derivatives such as: 2-methyl-benzene-1,4-diamine, benzene-1,4-diamine, 1-(2,5-diamino-phenyl)-ethanol, 2-(2,5-diamino-phenyl)-ethanol, N-(2-methoxyethyl)benzene-1,4-diamine, 2-[(4-amino-phenyl)-(2-hydroxy-ethyl)-amino]-ethanol, and 1-(2,5-diaminophenyl)ethane-1,2-diol;

25

p-aminophenol derivatives such as 4-amino-phenol, 4-methylamino-phenol, 4-amino-3-methyl-phenol, 4-amino-2-methoxymethyl-phenol, and 1-(5-amino-2-hydroxy-phenyl)-ethane-1,2-diol;

30

o-aminophenol derivatives such as: 2-amino-phenol, 2-amino-5-methyl-phenol, 2-amino-6-methyl-phenol, N-(4-amino-3-hydroxy-phenyl)-acetamide, and 2-amino-4-methyl-phenol; and

5 heterocyclic derivatives such as: pyrimidine-2,4,5,6-tetramine, 1-methyl-1H-pyrazole-4,5-diamine, 2-(4,5-diamino-1H-pyrazol-1-yl)ethanol, 1-(4-methylbenzyl)-1H-pyrazole-4,5-diamine, 1-(benzyl)-1H-pyrazole-4,5-diamine, and N<sup>2</sup>,N<sup>2</sup>-dimethyl-pyridine-2,5-diamine.

10 Preferred couplers include:

phenols, resorcinol and naphthol derivatives such as: naphthalene-1,7-diol, benzene-1,3-diol, 4-chlorobenzene-1,3-diol, naphthalen-1-ol, 2-methyl-naphthalen-1-ol, naphthalene-1,5-diol, naphthalene-2,7-diol, 15 benzene-1,4-diol, 2-methyl-benzene-1,3-diol, and 2-isopropyl-5-methylphenol;

m-phenylenediamines such as: benzene-1,3-diamine, 2-(2,4-diamino-phenoxy)-ethanol, 4-{3-[(2,4-diaminophenyl)oxy]propoxy}benzene-1,3-diamine, 2-(3-amino-4-methoxy-phenylamino)-ethanol, 2-[2,4-diamino-5- 20 (2-hydroxy-ethoxy)-phenoxy]-ethanol, and 3-(2,4-diamino-phenoxy)-propan-1-ol;

m-aminophenols such as: 3-amino-phenol, 5-amino-2-methyl-phenol, 5-(2-hydroxy-ethylamino)-2-methyl-phenol, and 3-amino-2-methyl- 25 phenol; and

heterocyclic derivatives such as: 3,4-dihydro-2H-1,4-benzoxazin-6-ol, 4-methyl-2-phenyl-2,4-dihydro-3H-pyrazol-3-one, 1,3-benzodioxol-5-ol, 1,3-benzodioxol-5-amine, 1H-indol-4-ol, 1H-indole-5,6-diol, 30 1H-indol-7-ol, 1H-indol-5-ol, 1H-indol-6-ol, 1H-indole-2,3-dione, pyridine-2,6-diamine, and 2-aminopyridin-3-ol.

Most preferred primary intermediates include:

5 p-phenylenediamine derivatives such as: 2-methyl-benzene-1,4-diamine, benzene-1,4-diamine, 2-(2,5-diamino-phenyl)-ethanol, 1-(2,5-diamino-phenyl)-ethanol, and 2-[(4-amino-phenyl)-(2-hydroxy-ethyl)-amino]-ethanol;

10 p-aminophenol derivatives such as: 4-amino-phenol, 4-methylamino-phenol, 4-amino-3-methyl-phenol, and 1-(5-amino-2-hydroxy-phenyl)-ethane-1,2-diol;

15 o-aminophenols such as: 2-amino-phenol, 2-amino-5-methyl-phenol, 2-amino-6-methyl-phenol, and N-(4-amino-3-hydroxy-phenyl)-acetamide; and

heterocyclic derivatives such as: pyrimidine-2,4,5,6-tetramine, 2-(4,5-diamino-1H-pyrazol-1-yl)ethanol, 1-(4-methylbenzyl)-1H-pyrazole-4,5-diamine, and 1-(benzyl)-1H-pyrazole-4,5-diamine.

20 Most preferred couplers include:

phenols, resorcinol and naphthol derivatives such as: benzene-1,3-diol, 4-chlorobenzene-1,3-diol, naphthalen-1-ol, 2-methyl-naphthalen-1-ol, and 2-methyl-benzene-1,3-diol;

25 m-phenylenediamine such as: 2-(2,4-diamino-phenoxy)-ethanol, 2-(3-amino-4-methoxy-phenylamino)-ethanol, 2-[2,4-diamino-5-(2-hydroxy-ethoxy)-phenoxy]-ethanol, and 3-(2,4-diamino-phenoxy)-propan-1-ol;

30 m-aminophenols such as: 3-amino-phenol, 5-amino-2-methyl-phenol, 5-(2-hydroxy-ethylamino)-2-methyl-phenol, and 3-amino-2-methyl-phenol; and

heterocyclic derivatives such as: 3,4-dihydro-2H-1,4-benzoxazin-6-ol, 4-methyl-2-phenyl-2,4-dihydro-3H-pyrazol-3-one, 1H-indol-6-ol, and 2-aminopyridin-3-ol.

5 Understandably, the coupler compounds and the primary intermediate compounds, including the novel compounds of the invention, in so far as they are bases, can be used as free bases or in the form of their physiologically compatible salts with organic or inorganic acids, such as hydrochloric, citric, acetic, tartaric, or sulfuric acids, or, in so far as they have  
10 aromatic OH groups, in the form of their salts with bases, such as alkali phenolates.

The total amount of dye precursors (e.g., primary intermediate and coupler compounds, including the novel compounds of this invention) in  
15 the hair dyeing compositions of this invention is generally from about 0.002 to about 20, preferably from about 0.04 to about 10, and most preferably from about 0.1 to about 7.0 weight percent, based on the total weight of the hair dyeing composition. The primary intermediate and coupler compounds are generally used in molar equivalent amounts. However, it is possible to use  
20 the primary intermediate compounds in either excess or deficiency, i.e., a molar ratio of primary intermediate to coupler generally ranging from about 5:1 to about 1:5.

The hair dyeing compositions of this invention will contain the  
25 primary intermediate of this invention in an effective dyeing amount, generally in an amount of from about 0.001 to about 10 weight percent by weight of the hair dye composition, preferably from about 0.01 to about 5.0 weight percent. Other primary intermediates, when present, are typically present in an amount such that in aggregate the concentration of primary intermediates in the  
30 composition is from about 0.002 to about 10 weight percent, preferably from about 0.01 to about 5.0 weight percent. The coupler(s) are present in an effective dyeing concentration, generally an amount of from about 0.001 to

about 10.0 weight percent by weight of the hair dye composition, preferably from about 0.01 to about 5.0 weight percent. The remainder of the hair dye composition comprises a carrier or vehicle for the couplers and primary intermediates, and comprises various adjuvants as described below.

5

Any suitable carrier or vehicle, generally an aqueous or hydroalcoholic solution, can be employed, preferably an aqueous solution. The carrier or vehicle will generally comprise more than 80 weight percent of the hair dye composition, typically 90 to 99 weight percent, preferably 94 to 99 weight percent. The hair coloring compositions of this invention may contain as adjuvants one or more cationic, anionic, amphoteric, or zwitterionic surface active agents, perfumes, antioxidants such as ascorbic acid, thioglycolic acid or sodium sulfite, chelating and sequestering agents such as EDTA, thickening agents, alkalizing or acidifying agents, solvents, diluents, inerts, dispersing agents, penetrating agents, defoamers, enzymes, and other dye agents (e.g., synthetic direct and natural dyes). These adjuvants are cosmetic additive ingredients commonly used in compositions for coloring hair.

10  
15

The hair dye compositions of the present invention are used by admixing them with a suitable oxidant, which reacts with the hair dye precursors to develop the hair dye. Any suitable oxidizing agent can be employed in the hair dye product compositions of this invention, particularly hydrogen peroxide ( $H_2O_2$ ) or precursors therefor. Also suitable are urea peroxide, the alkali metal salts of persulfate, perborate, and percarbonate, especially the sodium salt, and melamine peroxide. The oxidant is usually provided in an aqueous composition generally referred to as the developer composition, which normally is provided as a separate component of the finished hair dye product and present in a separate container. The developer composition may also contain, to the extent compatible, various ingredients needed to form the developer composition, i.e., peroxide stabilizers, foam formers, etc., and may incorporate one or more of the adjuvants referred to above, e.g., surface active agents, thickeners, pH modifiers, etc. Upon mixing the hair coloring composition and the developer composition to form a

20

25

30

hair dye product composition, the adjuvants are provided in the hair dye product composition as it is applied to the hair to achieve desired product attributes, e.g., pH, viscosity, rheology, etc.

5                   The form of the hair dye product compositions according to the invention can be, for example, a solution, especially an aqueous or aqueous-alcoholic solution. However, the form that is particularly preferred is a thick liquid, cream, gel or an emulsion whose composition is a mixture of the dye ingredients with the conventional cosmetic additive ingredients suitable for the  
10                   particular preparation.

                  Suitable conventional cosmetic additive ingredients useful in the hair dye and developer compositions, and hence in the hair dye product compositions of this invention are described below, and may be used to  
15                   obtain desired characteristics of the hair dye, developer and hair dye product compositions.

                  Solvents: In addition to water, solvents that can be used are lower alkanols (e.g., ethanol, propanol, isopropanol, benzyl alcohol); polyols (e.g., carbitols, propylene glycol, hexylene glycol, glycerin). See WO 98/27941 (section on  
20                   diluent) incorporated by reference. See also US 6027538 incorporated by reference. Under suitable processing, higher alcohols, such as C8 to C18 fatty alcohols, especially cetyl alcohol, are suitable organic solvents, provided they are first liquified by melting, typically at low temperature (50 to 80 °C), before  
25                   incorporation of other, usually lipophilic, materials.

                  The organic solvents are typically present in the hair dye compositions in an amount of from about 5 to about 30% by weight of the hair dye composition. Water is usually present in an amount of from about 5 to  
30                   about 90% by weight of the hair dye composition, preferably from about 15 to about 75% by weight and most preferably from about 30 to about 65% by weight.



Surfactants: These materials are from the classes of anionic, cationic, amphoteric (including zwitterionic surfactants) or nonionic surfactant compounds. (Cationic surfactants, generally included as hair conditioning materials, are considered separately below.) Suitable surfactants, other than

5 cationic surfactants, include fatty alcohol sulfates, ethoxylated fatty alcohol sulfates, alkylsulfonates, alkylbenzensulfonates, alkyltrimethylammonium salts, alkylbetaines, ethoxylated fatty alcohols, ethoxylated fatty acids, ethoxylated alkylphenols, block polymers of ethylene and/orpropylene glycol,

10 glycerol esters, phosphate esters, fatty acid alkanol amides and ethoxylated fatty acid esters, alkyl sulfates, ethoxylated alkyl sulfates, alkyl glyceryl ether sulfonates, methyl acyl taurates, acyl isethionates, alkyl ethoxy carboxylates, fatty acid mono- and diethanolamides. Especially useful are sodium and ammonium alkyl sulfates, sodium and ammonium ether sulfates having 1 to 3

15 ethylene oxide groups, and nonionic surfactants sold as Tergitols, e.g., C11-C15 Pareth-9, and Neodols, e.g., C12-C15 Pareth-3. They are included for various reasons, e.g., to assist in thickening, for forming emulsions, to help in wetting hair during application of the hair dye product composition, etc. Amphoteric surfactants include, for example, the asparagine derivatives as well

20 betaines, sultaines, glycinate and propionates having an alkyl or alkylamido group of from about 10 to about 20 carbon atoms. Typical amphoteric surfactants suitable for use in this invention include lauryl betaine, lauroamphoglycinate, lauroamphopropionate, lauryl sultaine, myristamidopropyl betaine, myristyl betaine, stearoamphopropylsulfonate,

25 cocamidoethyl betaine, cocamidopropyl betaine, cocoamphoglycinate, cocoamphocarboxypropionate, cocoamphocarboxyglycinate, cocobetaine, and cocoamphopropionate. Reference is made to WO 98/52523 published November 26, 1998 and WO 01/62221 published August 30, 2001, both incorporated herein by reference thereto.

30

The amount of surfactants in the hair dye compositions is normally from about 0.1% to 30% by weight, preferably 1% to 15% by weight.

Thickeners: Suitable thickeners include such as higher fatty alcohols, starches, cellulose derivatives, petrolatum, paraffin oil, fatty acids and anionic and nonionic polymeric thickeners based on polyacrylic and polyurethane polymers. Examples are hydroxyethyl cellulose, hydroxymethylcellulose and other cellulose derivatives, hydrophobically modified anionic polymers and nonionic polymers, particularly such polymers having both hydrophilic and hydrophobic moieties (i.e., amphiphilic polymers). Useful nonionic polymers include polyurethane derivatives such as PEG-150/stearyl alcohol/SDMI copolymer. Suitable polyether urethanes are Aculyn® 22, 44 and Aculyn® 46 polymers sold by Rohm & Haas. Other useful amphiphilic polymers are disclosed in US Pat. No. 6010541 incorporated by reference. See also WO 01/62221 mentioned above. Examples of anionic polymers that can be used as thickeners are acrylates copolymer, acrylates/ceteth-20 methacrylates copolymer, acrylates/ceteth-20 itaconate copolymer, and acrylates/beheneth-25 acrylates copolymers. In the case of the associative type of thickeners, e.g., Aculyns 22, 44 and 46, the polymer may be included in one of either the hair dye composition or the developer composition of the hair dye product and the surfactant material in the another. Thus, upon mixing of the hair dye and developer compositions, the requisite viscosity is obtained. The thickeners are provided in an amount to provide a suitably thick product as it is applied to the hair. Such products generally have a viscosity of from 1000 to 100000 cps, and often have a thixotropic rheology.

pH Modifying agents: Suitable materials that are used to adjust pH of the hair dye compositions include alkalizers such alkali metal and ammonium hydroxides and carbonates, especially sodium hydroxide and ammonium carbonate, ammonia, organic amines including methylethanolamine, aminomethylpropanol, mono-, di-, and triethanolamine, and acidulents such as inorganic and inorganic acids, for example phosphoric acid, acetic acid, ascorbic acid, citric acid or tartaric acid, hydrochloric acid, etc. See US patent 6027538 incorporated by reference.

Conditioners: Suitable materials include silicones and silicone derivatives; hydrocarbon oils; monomeric quaternary compounds, and quaternized polymers. Monomeric quaternary compounds are typically cationic compounds, but may also include betaines and other amphoteric and zwitterionic materials that provide a conditioning effect. Suitable monomeric quaternary compounds include behentrialkonium chloride, behentrimonium chloride, benzalkonium bromide or chloride, benzyl triethyl ammonium chloride, bis-hydroxyethyl tallowmonium chloride, C12-18 dialkyldimonium chloride, cetalkonium chloride, ceteartrimonium bromide and chloride, cetrimonium bromide, chloride and methosulfate, cetylpyridonium chloride, cocamidopropyl ethyldimonium ethosulfate, cocamidopropyl ethosulfate, coco-ethyldimonium ethosulfate, cocotrimonium chloride and ethosulfate, dibehenyl dimonium chloride, dicetyldimonium chloride, dicocodimonium chloride, dilauryl dimonium chloride, disoydimonium chloride, ditallowdimonium chloride, hydrogenated tallow trimonium chloride, hydroxyethyl cetyl dimonium chloride, myristalkonium chloride, olealkonium chloride, soyethomonium ethosulfate, soytrimonium chloride, stearalkonium chloride, and many other compounds. See WO 98/27941 incorporated by reference. Quaternized polymers are typically cationic polymers, but may also include amphoteric and zwitterionic polymers. Useful polymers are exemplified by polyquaternium-4, polyquaternium-6, polyquaternium-7, polyquaternium-8, polyquaternium-9, polyquaternium-10, polyquaternium-22, polyquaternium-32, polyquaternium-39, polyquaternium-44 and polyquaternium-47. Silicones suitable to condition hair are dimethicone, amodimethicone, dimethicone copolyol and dimethiconol. See also WO 99/34770 published July 15, 1999, incorporated by reference, for suitable silicones. Suitable hydrocarbon oils would include mineral oil.

Conditioners are usually present in the hair dye composition in an amount of from about 0.01 to about 5% by weight of the composition.

Direct Dyes: The hair dyeing compositions according to the invention can also contain compatible direct dyes including Disperse Black 9, HC Yellow 2, HC Yellow 4, HC Yellow 15, 4-nitro-o-phenylenediamine, 2-amino-6-chloro-4-nitrophenol, HC Red 3, Disperse Violet 1, HC Blue 2, Disperse Blue 3, and Disperse Blue 377. These direct dyes can be contained in the hair coloring compositions of the invention in an amount of from about 0.05 to 4.0 percent by weight.

10 Natural ingredients: For example, proteins and protein derivatives, and plant materials such as aloe, chamomile and henna extracts.

Other adjuvants include polysaccharides, alkylpolyglycosides, buffers, chelating and sequestrant agents, antioxidants, and peroxide stabilizing agents as mentioned in WO 01/62221, etc.

The adjuvants referred to above but not specifically identified that are suitable are listed in the International Cosmetics Ingredient Dictionary and Handbook, (Eighth Edition) published by The Cosmetics, Toiletry, and Fragrance Association, incorporated by reference. In particular reference is made to Volume 2, Section 3 (Chemical Classes) and Section 4 (Functions) are useful in identifying a specific adjuvant to achieve a particular purpose or multipurpose.

25 The above-mentioned conventional cosmetic ingredients are used in amounts suitable for their functional purposes. For example, the surfactants used as wetting agents, associative agents, and emulsifiers are generally present in concentrations of from about 0.1 to 30 percent by weight, the thickeners are useful in an amount of from about 0.1 to 25 percent by weight, and the hair care functional materials are typically used in concentrations of from about 0.01 to 5.0 percent by weight.

The hair dyeing product composition as it is applied to the hair, i.e., after mixing the hair dye composition according to the invention and the developer, can be weakly acidic, neutral or alkaline according to their composition. The hair dye compositions can have pH values of from about 6 to 11.5, preferably from about 6.8 to about 10, and especially from about 8 to about 10. The pH of the developer composition is typically acidic, and generally the pH is from about 2.5 to about 6.5, usually about 3 to 5. The pH of the hair dye and developer compositions is adjusted using a pH modifier as mentioned above.

10

In order to use the hair coloring composition for dyeing hair, the above-described hair coloring compositions according to the invention are mixed with an oxidizing agent immediately prior to use and a sufficient amount of the mixture is applied to the hair, according to the hair abundance, generally from about 60 to 200 grams. Some of the adjuvants listed above (e.g., thickeners, conditioners, etc.) can be provided in the dye composition or the developer, or both, depending on the nature of the ingredients, possible interactions, etc., as is well known in the art.

20

Typically, hydrogen peroxide, or its addition compounds with urea, melamine, sodium borate or sodium carbonate, can be used in the form of a 3 to 12 percent, preferably 6 percent, aqueous solution as the oxidizing agent for developing the hair dye. Oxygen can also be used as the oxidizing agent. If a 6 percent hydrogen peroxide solution is used as oxidizing agent, the weight ratio of hair coloring composition and developer composition is 5:1 to 1:5, but preferably 1:1. In general, the hair dyeing composition comprising primary intermediate(s) and coupler(s), including at least one of the compounds of formula (1), is prepared and then, at the time of use, the oxidizing agents, such as  $H_2O_2$ , contained in a developer composition is admixed therewith until an essentially homogenous composition is obtained, which is applied shortly after preparation to the hair to be dyed and permitted to remain in contact with the hair for a dyeing effective amount of time. The

30

10052741.011302

5 mixture of the oxidizing agent and the dye composition of the invention (i.e., the hair dye product composition) is allowed to act on the hair for about 2 to about 60 minutes, preferably about 15 to 45, especially about 30 minutes, at about 15 to 50°C, the hair is rinsed with water, and dried. If necessary, it is washed with a shampoo and rinsed, e.g., with water or a weakly acidic solution, such as a citric acid or tartaric acid solution. Subsequently the hair is dried. Optionally, a separate conditioning product may also be provided.

10 Together the hair dye composition of the present invention comprising the hair dye primary intermediate (1) and the developer composition comprising the oxidizing agent form a system for dyeing hair. This system may be provided as a kit comprising in a single package separate containers of the hair dye composition, the developer, the optional conditioner or other hair treatment product, and instructions for use.

15 Especially useful primary intermediates of formula (1) of this invention will provide hair coloring compositions having outstanding color fastness, especially light fastness, fastness to washing and fastness to rubbing.

### 20 Dyeing Example 1

25 The following composition shown in Table 1 can be used for dyeing Piedmont hair. 100 g of the dyeing composition is mixed with 100 g 20 volume hydrogen peroxide. The resulting mixture is applied to the hair and permitted to remain in contact with the hair for 30 minutes. The dyed hair is then shampooed, rinsed with water and dried. The ranges of ingredients set out in Table 1 are illustrative of useful concentrations of the recited materials in a hair dye product.

30

**TABLE 1**  
**Composition for Dyeing Hair**

Ingredients	Range (wt %)	Weight (%)
Cocamidopropyl betaine	0-25	17.00
Polyquaternium-22	0-7	5.00
Monoethanolamine <sup>1</sup>	0-15	2.00
Oleic Acid	2-22	0.75
Citric Acid	0-3	0.10
28% Ammonium hydroxide <sup>1</sup>	0-15	5.00
Behentrimonium chloride	1-5	0.50
Sodium sulfite	0-1	0.10
EDTA	0-1	0.10
Erythorbic acid	0-1	0.40
Ethoxydiglycol	1-10	3.50
C11-15 Pareth-9 (Tergitol 15-S-9)	0.5-5	1.00
C12-15 Pareth-3 (Neodol 25-3)	0.25-5	0.50
Isopropanol	2-10	4.00
Propylene glycol	1-12	2.00
p-phenylenediamine	0-5	1 mmole
N,N-Bis(hydroxyethyl)-p-phenylene diamine	0-5	1 mmole
3-Methyl-p-aminophenol	0-5	1 mmole
p-Aminophenol	0-5	1 mmole
Primary Intermediate of this invention	0.5-5	4 mmoles
5-Amino-2-Methyl Phenol <sup>2</sup>	0-5	3 mmoles
2,4-Diaminophenoxyethanol <sup>2</sup>	0-5	3 mmoles
M-Phenylenediamine <sup>2</sup>	0-5	1 mmole
Water	qs to 100.00	qs to 100.00

<sup>1</sup> In the aggregate, these ingredients are in the range of 2 to 15% by weight.

<sup>2</sup> At least one of these dye precursors is typically present.

Exemplary combinations of hair coloring components employing a novel primary intermediate of formula (1) of this invention are shown in Table 1 and in combinations C1 to C136 in Tables A through H. Reading down the columns in Tables A through H, the Xes demonstrate combinations of dyes that can be formulated according to the present invention. For example, in Combination No. C1 in Column 4 of Table A, a primary intermediate compound of formula (1) of this invention (Row 1 of Table A) where R<sub>1</sub> and R<sub>2</sub> are as defined before, can be combined with 2-aminophenol. Especially preferred as the primary intermediate compounds of

formula (1) in such combinations as identified in Table 1 and in Tables A through H are the following:

- 5                   4-amino-2-(1-phenylamino-ethyl)-phenol;  
                  4-amino-2-(1-piperidin-1-yl-ethyl)-phenol;  
                  4-amino-2-[1-(pyridin-3-yl-amino)-ethyl]-phenol;  
                  4-amino-2-(1-benzylamino-ethyl)-phenol;  
                  4-amino-2-[1-(4-methoxy)-phenylamino-ethyl]-phenol;  
                  4-amino-2-(1-amino-ethyl)-phenol;  
10               4-amino-2-(1-dimethylamino-ethyl)-phenol;  
                  4-amino-2-(1-morpholin-4-yl-ethyl)-phenol;  
                  4-amino-2-[1-(4-methoxy)-benzylamino-ethyl]-phenol;  
                  4-amino-2-(1-methylamino-ethyl)-phenol; and  
                  4-amino-2-[1-(2-hydroxyethylamino)-ethyl]-phenol.

15

20











Table A. Dye Combinations													
Structure	IUPAC Name	Name	C1	C2	C3	C4	C5	C6	C7	C8	C9	C10	C11
			X	X	X	X	X	X	X	X	X	X	X
	2-Methyl-benzene-1,4-diamine	p-Toluene-diamine											
	Benzene-1,4-diamine	p-Phenylene-diamine											
	2-[(4-Amino-phenyl)-(2-hydroxy-ethyl)-amino]-ethanol	N,N-Bis(2-hydroxyethyl)-p-phenylene-diamine											
	1-(2,5-Diamino-phenyl)-ethanol	1-Hydroxyethyl-p-phenylenediamine											
	4-Amino-3-methylphenol	3-Methyl-p-aminophenol											
	2-Amino-phenol	o-Aminophenol	X									X	
	Benzene-1,3-diol	Resorcinol		X								X	X

Table A (continued). Dye Combinations													
Structure	IUPAC Name	Name	C1	C2	C3	C4	C5	C6	C7	C8	C9	C10	C11
	2-Methyl-benzene-1,3-diol	2-Methyl-resorcinol			x								x
	Naphthalen-1-ol	1-Naphthol				x							
	2-Methyl-naphthalen-1-ol	2-Methyl-1-naphthol					x						
	2-(2,4-Diaminophenoxy)-ethanol	2,4-Diaminophenoxyethanol						x					
	Benzene-1,3-diamine	m-Phenylenediamine							x				
	3-Amino-phenol	m-Aminophenol								x			
	5-Amino-2-methylphenol	2-Hydroxy-4-aminotoluene									x		
	2-(4,5-Diaminopyrazol-1-yl)-ethanol	1-Hydroxyethyl-4,5-diamino-pyrazole											

















Table B. Dye Combinations																		
Structure	C12	C13	C14	C15	C16	C17	C18	C19	C20	C21	C22	C23	C24	C25	C26	C27	C28	C29
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
							X	X	X	X	X	X	X	X	X	X	X	X
																		
																		
																		
																		
							X									X		
	X	X	X	X	X	X		X									X	X

Table B (continued). Dye Combinations																		
Structure	C12	C13	C14	C15	C16	C17	C18	C19	C20	C21	C22	C23	C24	C25	C26	C27	C28	C29
									X								X	
	X									X								X
		X									X							
			X									X						
				X									X					
					X									X				
															X			
																		

Structure	C30	C31	C32	C33	C34	C35	C36	C37	C38	C39	C40	C41	C42	C43	C44	C45	C46	C47
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
	X	X	X	X	X													
						X	X	X	X	X	X	X	X	X	X	X	X	X
						X									X			
	X	X	X	X	X										X	X	X	X

Table C (continued). Dye Combinations









Structure	C30	C31	C32	C33	C34	C35	C36	C37	C38	C39	C40	C41	C42	C43	C44	C45	C46	C47
								X								X		
									X								X	
	X									X								X
		X									X							
			X									X						
				X									X					
					X									X				
																		









Table D. Dye Combinations																		
Structure	C48	C49	C50	C51	C52	C53	C54	C55	C56	C57	C58	C59	C60	C61	C62	C63	C64	C65
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
																		
	X	X	X	X														
																		
					X	X	X	X	X	X	X	X	X	X	X	X	X	X
																		
					X									X				
	X	X	X	X		X								X	X	X	X	X

Table D (continued). Dye Combinations









Structure	C48	C49	C50	C51	C52	C53	C54	C55	C56	C57	C58	C59	C60	C61	C62	C63	C64	C65
							X								X			
								X								X		
									X								X	
	X									X								X
		X									X							
			X									X						
				X									X					
																		











Table E. Dye Combinations																		
Structure	C66	C67	C68	C69	C70	C71	C72	C73	C74	C75	C76	C77	C78	C79	C80	C81	C82	C83
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
																		
																		
					X	X	X	X	X	X	X	X						
	X	X	X															
																		
				X									X					
	X	X	X		X								X	X	X	X	X	X

Table E (continued). Dye Combinations









Structure	C66	C67	C68	C69	C70	C71	C72	C73	C74	C75	C76	C77	C78	C79	C80	C81	C82	C83
						X								X				
							X								X			
								X								X		
									X								X	
	X									X								X
		X									X							
			X															
				X														X

Table F. Dye Combinations																		
Structure	C84	C85	C86	C87	C88	C89	C90	C91	C92	C93	C94	C95	C96	C97	C98	C99	C100	C101
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
			X	X	X	X	X	X	X	X	X							
</																		









Table F (continued). Dye Combinations																		
Structure	C84	C85	C86	C87	C88	C89	C90	C91	C92	C93	C94	C95	C96	C97	C98	C99	C100	C101
					X									X				
						X									X			
							X									X		
								X									X	
									X									X
	X									X								
		X									X							
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Table G. Dye Combinations																		
Structure	C102	C103	C104	C105	C106	C107	C108	C109	C110	C111	C112	C113	C114	C115	C116	C117	C118	C119
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
												X	X	X	X	X	X	X
	X	X																
			X	X	X	X	X	X	X	X	X							
			X															
			X															

Table G (continued). Dye Combinations


















Structure	C102	C103	C104	C105	C106	C107	C108	C109	C110	C111	C112	C113	C114	C115	C116	C117	C118	C119
					X								X					
						X								X				
							X								X			
								X								X		
									X								X	
	X									X								X
		X									X							
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Table H. Dye Combinations																	
Structure	C120	C121	C122	C123	C124	C125	C126	C127	C128	C129	C130	C131	C132	C133	C134	C135	C136
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
	X																
		X	X	X	X	X	X	X	X								

Table H (continued). Dye Combinations															
Structure	C120	C121	C122	C123	C124	C125	C126	C127	C128	C129	C130	C131	C132	C133	C134
			X								X				
				X								X			
					X								X		
						X								X	
							X								X
								X							X
									X						
	X														X
	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X



With the foregoing description of the invention, those skilled in  
5 the art will appreciate that modifications may be made to the invention without  
departing from the spirit thereof. Therefore, it is not intended that the scope of  
the invention be limited to the specific embodiments illustrated and described.